

THE BOTANICAL MAGAZINE TOKYO

VOL. 100 (1987)

Numbers 1057-1060, Pages 1-436

EDITORIAL BOARD

- | | |
|--|--|
| NOBORU HARA : | Editor-in-Chief, <i>Department of Biology, University of Tokyo, Komaba, Meguro-ku, Tokyo 153</i> |
| KUNIO IWATSUKI : | Editor, <i>Botanical Gardens, University of Tokyo, Hakusan, Bunkyo-ku, Tokyo 112</i> |
| MASAYUKI KATSUMI : | Editor, <i>Department of Biology, International Christian University, Mitaka, Tokyo 181</i> |
| ZENNOSKE IWATSUKI, <i>Hiroshima University</i> | MASAYUKI TAKAHASHI, <i>University of Tokyo</i> |
| SHOICHI KAWANO, <i>Kyoto University</i> | MASAKAZU TATEWAKI, <i>Hokkaido University</i> |
| ATSUSHI KOMAMINE, <i>Tohoku University</i> | MASAMITSU WADA, <i>Tokyo Metropolitan University</i> |
| TSUNEYOSHI KUROIWA, <i>University of Tokyo</i> | KYOJI YODA, <i>Osaka City University</i> |
| TATSUO OKU, <i>Kyushu University</i> | |

THE BOTANICAL SOCIETY OF JAPAN

CONTENTS

THE BOTANICAL MAGAZINE, TOKYO

Vol. 100

No. 1057 (March 1987)

| | |
|--|-----|
| MINAMIKAWA, T., N. MASUDA, A. KADOTA AND M. WADA: Effect of Sulfite on Compositional Changes of Cell Constituents in Germinating Spores of <i>Adiantum capillus-veneris</i> L. | 1 |
| TADA, T.: Size-Dependent Perennation Bud Formation and Its Effect on Seedling Establishment in <i>Veronicastrum sibiricum</i> ssp. <i>japonicum</i> | 9 |
| ITO, M.: Phylogenetic Systematics of the Nymphaeales | 17 |
| TAKENO, K. AND M. FURUYA: Sporophyte Formation in Experimentally-Induced Unisexual Female and Bisexual Gametophytes of <i>Lygodium japonicum</i> | 37 |
| YAMAMOTO, R.: Effect of Galactose on Auxin-Induced Cell Elongation in Oat Coleoptile Segments in Mannitol Solutions | 43 |
| WADA, M., H. SHIMIZU AND N. KONDO: A Model System to Study the Effect of SO ₂ on Plant Cells. II. Effect of Sulfite on Fern Spore Germination and Rhizoid Development | 51 |
| UENO, O. AND T. KOYAMA: Distribution and Evolution of C ₄ Syndrome in <i>Rhynchospora</i> (Rhynchosporae-Cyperaceae) | 63 |
| HOZUMI, K.: Analysis of Growth Curve of Stem Volume in Some Woody Species Using the <i>u-w</i> Diagram | 87 |
| SUGAWARA, T.: Chromosome Number of <i>Saruma</i> Oliver (Aristolochiaceae) | 99 |
| MASUZAWA, T.: A Comparison of the Photosynthetic Activity of <i>Polygonum weyrichii</i> var. <i>alpinum</i> under Field Conditions at the Timberline of Mt. Fuji and in the Laboratory | 103 |

No. 1058 (June 1987)

| | |
|--|-----|
| YAMAKURA, T.: An Empirical Approach to the Analysis of Forest Stratification I. Proposed Graphical Method Derived by Using an Empirical Distribution Function | 109 |
| YOKOHAMA, Y., J. TANAKA AND M. CHIHARA: Productivity of the <i>Ecklonia cava</i> Community in a Bay of Izu Peninsula on the Pacific Coast of Japan | 129 |
| YOSHITAMA, K., H. NISHINO, H. OZAWA, M. SAKATANI, Y. OKABE AND N. ISHIKURA: Distribution Pattern of Anthocyanidins and Anthocyanins in Polygonaceous Plants | 143 |
| HARA, Y. AND M. CHIHARA: Morphology, Ultrastructure and Taxonomy of the Raphidophycean Alga <i>Heterosigma akashiwo</i> | 151 |
| YOSHIDA, T. AND H. KAWAI: Taxonomic Study of the Genus <i>Myagropsis</i> (Cystoseiraceae, Phaeophyta) | 165 |
| PELLMYR, O.: Temporal Patterns of Ovule Allocation, Fruit Set, and Seed Predation in <i>Anemonopsis macrophylla</i> (Ranunculaceae) | 175 |
| TAKAHASHI, H.: A Comparative Floral and Pollination Biology of <i>Tricyrtis flava</i> Maxim., <i>T. nana</i> Yatabe and <i>T. ohsumiensis</i> Masamune (Liliaceae) | 185 |
| TERASAKA, O. AND T. NIITSU: Unequal Cell Division and Chromatin Differentiation in Pollen Grain Cells I. Centrifugal, Cold and Caffeine Treatments | 205 |
| ONO, K., Y. MURASAKI AND K. KAWAUCHI: Establishment and Characteristics of a Cell Suspension Culture from a Moss, <i>Atrichum undulatum</i> | 217 |

No. 1059 (September 1987)

| | |
|---|-----|
| SUZUKI, E. AND J. TSUKAHARA: Age Structure and Regeneration of Old Growth <i>Cryptomeria japonica</i> Forests on Yakushima Island | 223 |
| KOJIMA, Y., Y. NIINOMI, S. TSUBOI, T. HIYAMA AND H. SAKURAI: Destruction of Photosystem I Iron-Sulfur Centers of Spinach and <i>Anacystis nidulans</i> by Mercurials | 243 |
| YASUI, T., Y. ENDO AND H. OHASHI: Infrageneric Variation of the Low Molecular Weight Carbohydrate Composition of the Seeds of the Genus <i>Vicia</i> (Leguminosae) | 255 |
| TATEOKA, T. AND M. MICHIKAWA: <i>Agrostis mertensii</i> (Poaceae) in Hokkaido | 273 |
| KHAN, M.A.: Interspecies Interactions in Aquatic Hyphomycetes | 295 |
| KOHYAMA, T.: Stand Dynamics in a Primary Warm-Temperate Rain Forest Analyzed by the Diffusion Equation | 305 |
| AKIYAMA, H.: Studies on <i>Leucodon</i> (Leucodontaceae, Musci) and Related Genera in East Asia I. A Taxonomic Revision of <i>Leucodon</i> from Taiwan | 319 |

No. 1060 (December 1987)

| | |
|---|-----|
| SUGAWARA, T.: Taxonomic Studies of <i>Asarum</i> sensu lato - III. Comparative Floral Anatomy | 335 |
| SAKIO, H. AND T. MASUZAWA: Ecological Studies on the Timberline of Mt. Fuji II. Primary Productivity of <i>Alnus maximowiczii</i> Dwarf Forest | 349 |
| IMAICHI, R., D. DARNAEDI AND M. KATO: Adventitious Buds of <i>Dryopteris sparsa</i> Complex (Dryopteridaceae): Developmental Anatomy and Systematic Implication | 365 |
| MATSUDA, Y., A. MUSGRAVE, H. VAN DEN ENDE AND K. ROBERTS: Cell Walls of Algae in the Volvocales: Their Sensitivity to a Cell Wall Lytic Enzyme and Labeling with an Anti-Cell Wall Glycopeptide of <i>Chlamydomonas reinhardtii</i> | 373 |
| TAKAHASHI, H.: Pollen Morphology and Its Taxonomic Significance of the Monotropoideae (Ericaceae) | 385 |

INVITED ARTICLE

| | |
|---|-----|
| SUGIURA, M.: Structure and Function of the Tobacco Chloroplast Genome | 407 |
|---|-----|

AUTHOR INDEX

THE BOTANICAL MAGAZINE, TOKYO

Vol. 100

| | | | |
|----------------|----------|------------------|----------|
| AKIYAMA, H. | 319 | OHASHI, H. | 255 |
| CHIHARA, M. | 129, 151 | OKABE, Y. | 143 |
| DARNAEDI, D. | 365 | ONO, K. | 217 |
| ENDO, Y. | 255 | OZAWA, H. | 143 |
| FURUYA, M. | 37 | PELLMYR, O. | 175 |
| HARA, Y. | 151 | ROBERTS, K. | 373 |
| HIYAMA, T. | 243 | SAKATANI, M. | 143 |
| HOZUMI, K. | 87 | SAKIO, H. | 349 |
| IMAICHI, R. | 365 | SAKURAI, H. | 243 |
| ISHIKURA, N. | 143 | SHIMIZU, H. | 51 |
| ITO, M. | 17 | SUGAWARA, T. | 99, 335 |
| KADOTA, A. | 1 | SUGIURA, M. | 407 |
| KATO, M. | 365 | SUZUKI, E. | 223 |
| KAWAI, H. | 165 | TADA, T. | 9 |
| KAWAUCHI, K. | 217 | TAKAHASHI, H. | 185, 385 |
| KHAN, M.A. | 295 | TAKENO, K. | 37 |
| KOBYAMA, T. | 305 | TANAKA, J. | 129 |
| KOJIMA, Y. | 243 | TATEOKA, T. | 273 |
| KONDO, N. | 51 | TERASAKA, O. | 205 |
| KOYAMA, T. | 63 | TSUBOI, S. | 243 |
| MASUDA, N. | 1 | TSUKAHARA, J. | 223 |
| MASUZAWA, T. | 103, 349 | UENO, O. | 63 |
| MATSUDA, Y. | 373 | VAN DEN ENDE, H. | 373 |
| MICHIKAWA, M. | 273 | WADA, M. | 1, 51 |
| MINAMIKAWA, T. | 1 | YAMAKURA, T. | 109 |
| MURASAKI, Y. | 217 | YAMAMOTO, R. | 43 |
| MUSGRAVE, A. | 373 | YASUI, T. | 255 |
| NIINOMI, Y. | 243 | YOKOHAMA, Y. | 129 |
| NIITSU, T. | 205 | YOSHIDA, T. | 165 |
| NISHINO, H. | 143 | YOSHITAMA, K. | 143 |

Index of Key Words

| | | | | | |
|----------------------------------|---------|--|------------------------------------|-----|--|
| A | | | | | |
| <i>Adiantum capillus-veneris</i> | 1 | | Diffusion equation | 305 | |
| Adventitious bud | 365 | | Distribution pattern | 143 | |
| Age structure | 223 | | <i>Dryopteris sparsa</i> complex | 365 | |
| <i>Agrostis flaccida</i> | 273 | | E | | |
| <i>Agrostis mertensii</i> | 273 | | <i>Ecklonia cava</i> | 129 | |
| <i>Agrostis tateyamensis</i> | 273 | | Ecological distribution | 63 | |
| Air pollution | 51 | | F | | |
| <i>Alnus maximowiczii</i> | 349 | | <i>Fagus crenata</i> | 87 | |
| Amino acids | 1 | | Fern | 51 | |
| <i>Anacystis nidulans</i> | 243 | | Flora of Hokkaido | 273 | |
| <i>Anemonopsis</i> | 175 | | Floral anatomy | 335 | |
| Annual production | 349 | | Floral biology | 185 | |
| Antheridiogen | 37 | | Forest phytomass | 349 | |
| Anthocyanidins | 143 | | Forest regeneration | 223 | |
| Anthocyanins | 143 | | Forest stratification | 109 | |
| Aquatic hyphomycetes | 295 | | G | | |
| Aristolochiaceae | 99, 335 | | Galactose | 43 | |
| <i>Asarum</i> | 99, 335 | | Gap dynamics | 305 | |
| <i>Atrichum undulatum</i> | 217 | | Geographical distribution | 63 | |
| Autogamy | 185 | | Germination | 51 | |
| Auxin | 43 | | Germination (fern spore) | 1 | |
| <i>Avena sativa</i> | 43 | | Germination time | 9 | |
| B | | | Gibberellin | 37 | |
| <i>Bombus</i> | 185 | | Growth curve | 87 | |
| C | | | H | | |
| C ₄ syndrome | 63 | | <i>Heterosigma akashiwo</i> | 151 | |
| Carpel evolution | 175 | | <i>Heterosigma inlandica</i> | 151 | |
| Cell wall | 373 | | <i>Heterotropa</i> | 335 | |
| Cell wall synthesis | 43 | | I | | |
| Ceratophyllaceae | 17 | | Infraspecific variation | 385 | |
| <i>Chamaecyparis obtusa</i> | 87 | | Inhibitors of photosynthesis | 243 | |
| Chemotaxonomy | 255 | | Interspecies interactions | 295 | |
| <i>Chlamydomonas reinhardtii</i> | 373 | | Intron | 407 | |
| Chloroplast DNA | 407 | | Iron-sulfur centers | 243 | |
| Chromatin differentiation | 205 | | K | | |
| Chromosome number | 99 | | Kranz leaf anatomy | 63 | |
| Cladistic analysis | 17 | | L | | |
| Complete nucleotide sequence | 407 | | Leaf area index | 129 | |
| <i>Cryptomeria japonica</i> | 87, 223 | | Leaf-marking method | 129 | |
| Cyperaceae | 63 | | <i>Leucodon</i> | 319 | |
| <i>Cystophyllum</i> | 165 | | Low molecular weight carbohydrates | 255 | |
| Cystoseiraceae | 165 | | <i>Lygodium japonicum</i> | 37 | |
| Cytogeography | 273 | | Lytic enzyme | 373 | |
| D | | | | | |
| Dbh distribution | 305 | | | | |
| Demography | 9 | | | | |
| Developmental anatomy | 365 | | | | |
| Difference equation | 109 | | | | |

M

| | |
|-------------------|----------|
| Mercurials | 243 |
| Monotropoideae | 385 |
| Moss | 217 |
| Mt. Fuji | 103, 349 |
| <i>Myagropsis</i> | 165 |

N

| | |
|----------------|-----|
| Nelumbonaceae | 17 |
| Net production | 129 |
| New World | 385 |
| Nymphaeaceae | 17 |
| Nymphaeales | 17 |

O

| | |
|-----------------------------|-----|
| Oat coleoptile | 43 |
| Old World | 385 |
| <i>Olisthodiscus luteus</i> | 151 |

P

| | |
|----------------------------|----------|
| Perennation bud | 9 |
| Peristome structure | 319 |
| Phaeophyta | 165 |
| Photosynthesis | 103, 407 |
| Photosystem I | 243 |
| Phylogenetic systematics | 17 |
| Pollen grain | 205 |
| Pollen morphology | 385 |
| Polygonaceae | 143 |
| <i>Polygonum weyrichii</i> | 103 |
| Polyploid complex | 273 |
| Proanthocyanidin | 143 |
| Production structure | 349 |
| Profile diagram | 109 |
| Protonema | 217 |
| Protoplast | 217 |

R

| | |
|---------------------|-----|
| Ranunculaceae | 175 |
| Raphidophycean alga | 151 |
| Reserve lipid | 1 |
| Reserve protein | 1 |
| Rhizoid | 51 |
| <i>Rhynchospora</i> | 63 |

S

| | |
|------------------------|-------|
| <i>Saruma</i> | 99 |
| Seasonality | 295 |
| Seedling establishment | 9 |
| Seeds | 255 |
| Sex differentiation | 37 |
| SO ₂ | 51 |
| Spatial allocation | 175 |
| Spinach chloroplasts | 243 |
| Sporophyte formation | 37 |
| Stand basal area | 305 |
| Stored growth | 43 |
| Submarine forest | 129 |
| Sulfite | 1, 51 |
| Suspension culture | 217 |

T

| | |
|------------------------------|-------------------------|
| Taiwan | 319 |
| Taxonomy | 151, 165, 319, 335, 385 |
| Temperate coniferous forest | 223 |
| Temporal ovule allocation | 175 |
| Timberline | 103, 349 |
| Tobacco | 407 |
| <i>Trans</i> splicing | 407 |
| <i>Tricyrtis flava</i> | 185 |
| <i>Tricyrtis nana</i> | 185 |
| <i>Tricyrtis ohsumiensis</i> | 185 |

U

| | |
|--------------------|-----|
| <i>u-w</i> diagram | 87 |
| Ultrastructure | 151 |
| Unequal division | 205 |

V

| | |
|-------------------------------------|-----|
| Vegetative and reproductive success | 295 |
| Vegetative reproduction | 365 |
| <i>Veronicastrum sibiricum</i> | 9 |
| <i>Vicia</i> | 255 |
| Volvocales | 373 |

W

| | |
|----------------------------|-----|
| Warm-temperate rain forest | 305 |
| Winter mortality | 9 |

Y

| | |
|------------------|----------|
| Yakushima Island | 223, 305 |
|------------------|----------|

Acknowledgement to Reviewers for Vol. 100

The Editorial Board is grateful to the following reviewers for their cooperation in examining the manuscripts submitted to the Botanical Magazine, Tokyo, Vol. 100.

ABE, K.
ESASHI, Y.
FUJII, T.
FUJITA, N.
HARA, T.
HATANAKA, S.
HIROSE, T.
HOSON, T.
HOZUMI, K.
INOUE, H.
ISHIKURA, N.
ISHIZUKA, K.
ITO, M.
IWAKI, H.
KANAI, H.
KATO, K.
KATO, M.
KATO, S.
KAZAMA, H.
KIMURA, M.
KODAMA, T.
KURAISHI, S.
KUROIWA, S.
MARUTA, E.
MASUDA, M.
MIKI, H.
MINAMIKAWA, T.
MOROHASHI, M.
NIITSU, T.
NISHIMURA, M.

NISHINO, E.
OHASHI, H.
OHBA, H.
OHISHI, K.
OIKAWA, T.
OKADA, H.
OKUDA, S.
SAEKI, T.
SAHASHI, N.
SHIBAOKA, H.
SOHMA, K.
SOMA, K.
SUGANUMA, T.
SUGIYAMA, J.
SUZUKI, M.
TAGAWA, H.
TAMURA, M.
TANAKA, R.
TATEOKA, T.
TERAMOTO, E.
TOBE, H.
TOTSUKA, T.
TSUBAKI, K.
UMEZAKI, I.
WATANABE, K.
YAMADA, M.
YAMAMOTO, Y.
YAMAZAKI, T.
YOSHIDA, O.
YOSHIDA, S.

INSTRUCTIONS TO AUTHORS (Revised November, 1987)

1. The Botanical Magazine, Tokyo is an international journal dedicated to the dissemination of fundamental knowledge in all areas of plant sciences. Only original papers, including short communications, and invited review articles in English will be considered. Short communications should contain important new information and will be published in the earliest possible issue. Manuscripts are accepted for review with the understanding that the same work has not been and will not be published nor is presently being submitted elsewhere and that all persons listed as authors have given their approval for the submission of the paper. By submitting a manuscript, the authors agree that the copyright for their article will be transferred to the Botanical Society of Japan if and when the article is accepted for publication. The copyright covers the exclusive rights to reproduce and distribute the article, including reprints, photographic reproductions, microform and other reproductions of similar nature, and translations.
2. Manuscripts should be written in clear, concise and grammatical English. In addition to the original typewritten manuscript, two copies each of text material, tables and figures should be submitted. Photographic figures must be submitted in triplicate. Authors who are uncertain about proper English usage should have their manuscripts checked prior to submission by someone proficient in the English language.
3. Full papers are limited to 16 printed pages in length including tables, figures and abstract. Short communications are limited to four printed pages. With the approval of the Editorial Board, additional pages can be published only at the author's expense (2,000 yen per page).
4. If none of the authors is a member of the Botanical Society of Japan, the authors must agree to pay a charge of 2,000 yen for each printed page. In this case, the manuscript, if accepted, will be published only after commitment to pay the charge by the authors or the financial officer of their institution.
5. Preparation of Manuscript Copy :
 - A. Double-space everything. Use good-quality paper of 28×21 cm or A4 size. Each typewritten page usually consists of 25 lines, each line having approximately 65 characters (two and half typewritten pages should be approximately equivalent to one printed page).
 - B. Arrange the manuscript copy in the order listed below and number all pages consecutively, including tables and figures, in the upper right-hand corner. In addition, indicate the following items on a separate cover sheet: (1) name and address for mailing galley proofs, (2) running title with less than 40 characters, and (3) the number of tables and figures.
 - a. Title of the paper, authors' names and institutions and footnotes (abbreviations, symbols, etc.)
 - b. Abstract: This should be a lucid digest of the paper, not exceeding 200 words for a full paper, less than 50 words for a short communication.
 - c. Key words: Alphabetically arrange less than 6 words or phrases which identify the most important subjects covered by the paper.
 - d. Text: Preparation of the text with the headings of Introduction, Materials and Methods, Results, and Discussion is suggested. Short communications should follow the style of full papers, but the headings should be omitted.
 - e. References cited: References cited in the text should be arranged alphabetically according to author on a separate sheet. Each reference should be given in the following order: (1) authors' names, (2) year of publication, (3) title of the

paper, (4) name of the journal (or book), (5) volume and page. Draw a wavy line under the volume number. For a single book, the name of the publisher and its location should be indicated.

Examples :

Bassham, J.A. and M. Calvin. 1957. The Path of Carbon in Photosynthesis. Prentice-Hall, Englewood Cliffs, New Jersey.

Esau, E. 1945. Vascularization of the vegetative shoots in *Helianthus* and *Sambucus*. Amer. J. Bot. **32** : 18-29.

Torrey, J.Q. 1965. Physiological bases of organization and development in the root. In A. Lang, ed., Encyclopedia of Plant Physiology, vol. 15/1, pp. 1256-1327. Springer-Verlag, Berlin.

- f. Tables : In addition to the heading, a lucid legend to explain the meaning of the content without requiring reference to the text should be included below the table. With the approval of the Editorial Board, the folders of tables can be published only at the authors' expense.
- g. Legends for figures and plates : In addition to the heading, a lucid legend should explain the meaning of the figure without requiring reference to the text. These legends should be grouped on a separate sheet.
- h. Figures and plates : Plates submitted should be of a size less than 27×18 cm. On the back of each figure or plate, indicate the authors' names, the number of the figure and its orientation. If a scale is necessary for the plate, it should be indicated. With the approval of the Editorial Board, colored illustrations can be published only at the authors' expense.

- C. Text references should be made using the authors' names (for three or more authors use Doe *et al.*) followed by the year of publication.

Example :

Hall and Cocking (1966) reported that ———.

——— (Hall and Cocking, 1966).

- D. The place of insertion of tables and/or figures in the text should be indicated on the right-hand margin of the sheet.
 - E. Units of measurement should of the metric system. Use Arabic numerals. For the use of Latin names, consult the International Code of Botanical Nomenclature.
- 6. Manuscripts not conforming to the aforementioned rules will be returned to the authors.
 - 7. Whether or not the manuscript is accepted, amendment of the manuscript and the order of publication are to be decided by the Editorial Board.
 - 8. Galley proofs will be sent directly from the printing company to the person designated. The authors are responsible for reading the first galley proof. No change of the content of the manuscript is permitted on the galley proof. If the Editorial Board recognizes the necessity for a change, this change will be made at the authors' expense.
 - 9. The authors will receive free of charge 50 offprints without covers. Additional copies and covers can be obtained at cost.
 - 10. All manuscripts should be sent by registered mail to the following address :
Editorial Office
The Botanical Society of Japan
Toshin Building
Hongo 2-27-2, Bunkyo-ku, Tokyo 113, Japan
 - 11. The manuscripts of the papers published will not be returned to the authors. However, figures and plates will be returned upon request by the authors after the paper has been published.